
Russian Federation


Total population (2016): 143 439 832 > Population aged 15 years and older (15+): 83% > Population in urban areas: 75% > Income group (World Bank): Upper-middle income

ALCOHOL CONSUMPTION: LEVELS AND PATTERNS

Recorded alcohol per capita (15+) consumption, 1961-2016



Recorded alcohol per capita (15+) consumption (in litres of pure alcohol) by type of alcoholic beverage, 2016 or latest year available



Alcohol per capita (15+) consumption (in litres of pure alcohol)

	20	10*	20	16*	
Recorded	11.1		8.1		
Unrecorded	4.7		3.6		
Total**	15.8		11.7		
Total males / females	26.1	7.3	18.7	5.8	
WHO European Region	11.2		9	9.8	

 $^{^{\}star}$ Three-year averages of recorded and unrecorded for 2009–2011 and 2015–2017; ** adjusted for tourist consumption.

Total alcohol per capita (15+) consumption, drinkers only (in litres of pure alcohol), 2016

Beer

Wine

Spirits Other All

	Litres
Males (15+)	30.5
Females (15+)	10.5
Both sexes (15+)	20.1

Prevalence of heavy episodic drinking* (%), 2016

	Population (15+ years)	Drinkers only (15+ years)	Population (15–19 years)	Drinkers only (15–19 years)
Males	48.4	78.8	37.9	85.1
Females	24.2	43.7	18.8	55.1
Both sexes	35.2	60.6	28.6	72.4

^{*} Consumed at least 60 grams or more of pure alcohol on at least one occasion in the past 30 days.

Abstainers (%), 2016

	Males	Females	Both sexes
Lifetime abstainers (15+)	24.0	30.0	27.2
Former drinkers* (15+)	14.7	14.6	14.6
Abstainers (15+), past 12 months	38.6	44.6	41.9

 $^{^{\}ast}$ Persons who used to drink alcoholic beverages but have not done so in the past 12 months.

HEALTH CONSEQUENCES: MORTALITY AND MORBIDITY

Age-standardized death rates (ASDR) and alcohol-attributable fractions (AAF), 2016

	ASI	OR*	AAF (%)		AAD** (Number)
Liver cirrhosis, males / females	40.3	18.4	51.7	61.8	22 850
Road traffic injuries, males / females	32.3	9.6	46.4	45.8	11 537
Cancer, males / females	298.4	153.1	10.1	5.5	28 702

^{*}Per 100 000 population (15+); **alcohol-attributable deaths, both sexes.

Years of life lost (YLL) score*, 2016 LEAST < 1 2 3 4 6 > MOST

Prevalence of alcohol use disorders and alcohol dependence (%), 2016^{*}

	Alcohol use disorders**	Alcohol dependence
Males	36.9	16.5
Females	7.4	3.3
Both sexes	20.9	9.3
WHO European Region	8.8	3.7

^{* 12-}month prevalence estimates (15+); **including alcohol dependence and harmful use of alcohol.

POLICIES AND INTERVENTIONS

Written national policy (adopted/revised) / National action plan	Yes (2009/2015) / No
Excise tax on beer / wine / spirits	Yes / Yes / Yes
National legal minimum age for off-premise sales of alcoholic beverages (beer / wine / spirits)	18 / 18 / 18
National legal minimum age for on-premise sales of alcoholic beverages (beer / wine / spirits)	18 / 18 / 18
Restrictions for on-/off-premise sales of alcoholic beverages (any): Hours, days / places, density Specific events / intoxicated persons / petrol stations	Yes, No / Yes, No Yes / No / Yes

National maximum legal blood alcohol concentration (BAC) when driving a vehicle (general / young / professional), in %	0.035 / 0.035 / 0.035
Legally binding regulations on alcohol advertising / product placement (any)	Yes / Yes
Legally binding regulations on alcohol sponsorship / sales promotion (any)	Yes / Yes
Legally required health warning labels on alcohol advertisements / containers (any)	Yes / Yes
National government support for community action (any)	Yes
National monitoring system(s) (any)	Yes

^{*} Based on alcohol-attributable years of life lost.